arXiv:1604.04973v1 [math.GR] 18 Apr 2016

Factorization numbers of finite rank 3 abelian p-groups

Marius Tărnăuceanu

March 22, 2016

Abstract

In this short note we give a formula for the factorization number $F_2(G)$ of a finite rank 3 abelian p-group G. This extends a result in our previous work [9].

MSC (2010): Primary 20D40; Secondary 20D60.
Key words: factorization number, subgroup commutativity degree, Möbius function, finite abelian p-group.

1 Introduction

Let G be a finite group, $L(G)$ be the subgroup lattice of G and H, K be two subgroups of G. If $G = HK$, then G is said to be *factorized* by H and K and the expression $G = HK$ is said to be a *factorization* of G. Denote by $F_2(G)$ the *factorization number* of G, that is the number of all factorizations of G.

The starting point for our discussion is given by the papers [6, 1], where $F_2(G)$ has been computed for certain classes of finite groups. Then, in [9], we have obtained explicit formulas for the factorization numbers of an elementary abelian p-group and of a rank 2 abelian p-group. These are based on the connection between $F_2(G)$ and the subgroup commutativity degree $sd(G)$ of G (see [7, 8]), namely

$$sd(G) = \frac{1}{|L(G)|^2} \sum_{H \leq G} F_2(H).$$

Obviously, by applying the well-known Möbius inversion formula to the above equality, one obtains

$$F_2(G) = \sum_{H \leq G} sd(H)|L(H)|^2 \mu(H, G). \tag{1}$$

In particular, if G is abelian, then we have $sd(H) = 1$ for all $H \in L(G)$, and consequently

$$F_2(G) = \sum_{H \leq G} |L(H)|^2 \mu(H, G) = \sum_{H \leq G} |L(G/H)|^2 \mu(H). \tag{2}$$

This formula will be used in the following to calculate the factorization number of a rank 3 abelian p-group.

First of all, we recall a theorem due to P. Hall [2] (see also [4]), that permits us to compute explicitly the Möbius function of a finite p-group.

Theorem 1. *Let G be a finite p-group of order p^n. Then $\mu(G) = 0$ unless G is elementary abelian, in which case we have $\mu(G) = (-1)^n p^{\binom{n}{2}}$.*

We also need to know the total number of subgroups of a finite rank 3 abelian p-group. It has been determined by using different methods in [3, 5, 10].

Theorem 2. *The total number of subgroups of $\mathbb{Z}_{p^{\lambda_1}} \times \mathbb{Z}_{p^{\lambda_2}} \times \mathbb{Z}_{p^{\lambda_3}}$, where $\lambda_1 \geq \lambda_2 \geq \lambda_3 \geq 1$, is*

$$f(\lambda_1, \lambda_2, \lambda_3) = \frac{A}{(p^2-1)^2(p-1)}, \tag{3}$$

where

$$\begin{aligned} A =& (\lambda_3+1)(\lambda_1-\lambda_2+1)p^{\lambda_2+\lambda_3+5} + 2(\lambda_3+1)p^{\lambda_2+\lambda_3+4} - 2(\lambda_3+1)(\lambda_1-\lambda_2)p^{\lambda_2+\lambda_3+3} \\ & - 2(\lambda_3+1)p^{\lambda_2+\lambda_3+2} + (\lambda_3+1)(\lambda_1-\lambda_2-1)p^{\lambda_2+\lambda_3+1} - (\lambda_1+\lambda_2-\lambda_3+3)p^{2\lambda_3+4} \\ & - 2p^{2\lambda_3+3} + (\lambda_1+\lambda_2-\lambda_3-1)p^{2\lambda_3+2} + (\lambda_1+\lambda_2+\lambda_3+5)p^2 + 2p - (\lambda_1+\lambda_2+\lambda_3+1). \end{aligned}$$

We are now able to give the main result of our note.

Theorem 3. *The following equality holds*

$$\begin{aligned}F_2(\mathbb{Z}_{p^{\lambda_1}}\times\mathbb{Z}_{p^{\lambda_2}}\times\mathbb{Z}_{p^{\lambda_3}})=&-p^3f^2(\lambda_1-1,\lambda_2-1,\lambda_3-1)+p(f^2(\lambda_1-1,\lambda_2-1,\lambda_3)\\&+pf^2(\lambda_1-1,\lambda_2,\lambda_3-1)+p^2f^2(\lambda_1,\lambda_2-1,\lambda_3-1))\\&-(f^2(\lambda_1-1,\lambda_2,\lambda_3)+pf^2(\lambda_1,\lambda_2-1,\lambda_3)\\&+p^2f^2(\lambda_1,\lambda_2,\lambda_3-1))+f^2(\lambda_1,\lambda_2,\lambda_3),\end{aligned}$$

where the quantities $f(\lambda_1,\lambda_2,\lambda_3)$ *are given by* (3).

We remark that for $\lambda_3=0$ the above equality leads to the formula in Theorem 3 of [9]. It simplifies in the particular case $\lambda_1=\lambda_2=\lambda_3=\lambda$.

Corollary 4. *We have*

$$\begin{aligned}F_2(\mathbb{Z}_{p^\lambda}\times\mathbb{Z}_{p^\lambda}\times\mathbb{Z}_{p^\lambda})=&-p^3f^2(\lambda-1,\lambda-1,\lambda-1)+p(1+p+p^2)f^2(\lambda,\lambda-1,\lambda-1)\\&-(1+p+p^2)f^2(\lambda,\lambda,\lambda-1)+f^2(\lambda,\lambda,\lambda).\end{aligned}$$

However, even in this case, an explicit formula for the factorization number is too difficult to be written, but we can do it for small values of $\lambda_1,\lambda_2,\lambda_3$.

Examples.

a) $F_2(\mathbb{Z}_{p^3}\times\mathbb{Z}_{p^2}\times\mathbb{Z}_p)=9p^6+15p^5+21p^4+16p^3+20p^2+11p+13.$

b) $F_2(\mathbb{Z}_{p^2}\times\mathbb{Z}_{p^2}\times\mathbb{Z}_{p^2})=5p^8+7p^7+16p^6+15p^5+21p^4+16p^3+20p^2+11p+13.$

2 Proof of Theorem 3

It is well-known that $G=\mathbb{Z}_{p^{\lambda_1}}\times\mathbb{Z}_{p^{\lambda_2}}\times\mathbb{Z}_{p^{\lambda_3}}$ has a unique elementary abelian subgroup of order p^3, say M, and that

$$G/M\cong\Phi(G)\cong\mathbb{Z}_{p^{\lambda_1-1}}\times\mathbb{Z}_{p^{\lambda_2-1}}\times\mathbb{Z}_{p^{\lambda_3-1}},$$

where $\Phi(G)$ is the Frattini subgroup of G. Moreover, all elementary abelian subgroups of G are contained in M. Denote by M_i, $i=1,2,...,p^2+p+1$, the subgroups of order p and by M_i', $i=1,2,...,p^2+p+1$, the subgroups of order p^2 in M. Then every quotient G/M_i is isomorphic to a maximal subgroup of G, which are: p^2 of type $(\lambda_1,\lambda_2,\lambda_3-1)$, p of type $(\lambda_1,\lambda_2-1,\lambda_3)$, and

1 of type $(\lambda_1 - 1, \lambda_2, \lambda_3)$. Similarly, every quotient G/M_i' is isomorphic to a subgroup of index p^2 of G that contains $\Phi(G)$, and these are: p^2 of type $(\lambda_1, \lambda_2 - 1, \lambda_3 - 1)$, p of type $(\lambda_1 - 1, \lambda_2, \lambda_3 - 1)$, and 1 of type $(\lambda_1 - 1, \lambda_2 - 1, \lambda_3)$. In this way the equality (2) becomes

$$F_2(G) = |L(G/M)|^2\mu(M) + \sum_{i=1}^{p^2+p+1} |L(G/M_i)|^2\mu(M_i) + \sum_{i=1}^{p^2+p+1} |L(G/M_i')|^2\mu(M_i') + |L(G)|^2\mu(1),$$

in view of Theorem 1. Since we have $\mu(M) = \mu(\mathbb{Z}_p^3) = -p^3$, $\mu(M_i) = \mu(\mathbb{Z}_p) = -1$, $\mu(M_i') = \mu(\mathbb{Z}_p^2) = p$, $\forall\, i = 1, 2, ..., p^2 + p + 1$, and $\mu(1) = 1$, one obtains

$$\begin{aligned} F_2(G) = & -p^3|L(\mathbb{Z}_{p^{\lambda_1-1}} \times \mathbb{Z}_{p^{\lambda_2-1}} \times \mathbb{Z}_{p^{\lambda_3-1}})|^2 + p(|L(\mathbb{Z}_{p^{\lambda_1-1}} \times \mathbb{Z}_{p^{\lambda_2-1}} \times \mathbb{Z}_{p^{\lambda_3}})|^2 \\ & + p|L(\mathbb{Z}_{p^{\lambda_1-1}} \times \mathbb{Z}_{p^{\lambda_2}} \times \mathbb{Z}_{p^{\lambda_3-1}})|^2 + p^2|L(\mathbb{Z}_{p^{\lambda_1}} \times \mathbb{Z}_{p^{\lambda_2-1}} \times \mathbb{Z}_{p^{\lambda_3-1}})|^2) \\ & - (|L(\mathbb{Z}_{p^{\lambda_1-1}} \times \mathbb{Z}_{p^{\lambda_2}} \times \mathbb{Z}_{p^{\lambda_3}})|^2 + p|L(\mathbb{Z}_{p^{\lambda_1}} \times \mathbb{Z}_{p^{\lambda_2-1}} \times \mathbb{Z}_{p^{\lambda_3}})|^2 \\ & + p^2|L(\mathbb{Z}_{p^{\lambda_1}} \times \mathbb{Z}_{p^{\lambda_2}} \times \mathbb{Z}_{p^{\lambda_3-1}})|^2) + |L(\mathbb{Z}_{p^{\lambda_1}} \times \mathbb{Z}_{p^{\lambda_2}} \times \mathbb{Z}_{p^{\lambda_3}})|^2. \end{aligned}$$

Under the notation of Theorem 2 this leads to the desired formula. ∎

References

[1] M. Farrokhi D.G., *Factorization numbers of finite abelian groups*, Int. J. Group Theory **2** (2013), 1-8.

[2] P. Hall, *A contribution to the theory of groups of prime-power order*, Proc. London Math. Soc. **36** (1933), 29-95.

[3] M. Hampejs and L. Tóth, *On the subgroups of finite abelian groups of rank three*, Annales Univ. Sci. Budapest., Sect. Comp., **39** (2013), 111-124.

[4] T. Hawkes, I.M. Isaacs and M. Özaydin, *On the Möbius function of a finite group*, Rocky Mountain J. Math. **19** (1989), 1003-1033.

[5] J.M. Oh, *An explicit formula for the number of subgroups of a finite abelian p-group up to rank* 3, Commun. Korean Math. Soc. **28** (2013), 649-667.

[6] F. Saeedi and M. Farrokhi D.G., *Factorization numbers of some finite groups*, Glasgow Math. J. **54** (2012), 345-354.

[7] M. Tărnăuceanu, *Subgroup commutativity degrees of finite groups*, J. Algebra **321** (2009), 2508-2520.

[8] M. Tărnăuceanu, *Addendum to "Subgroup commutativity degrees of finite groups"*, J. Algebra **337** (2011), 363-368.

[9] M. Tărnăuceanu, *On the factorization numbers of some finite p-groups*, accepted for publication in Ars Combin., arXiv:1502.04778.

[10] M. Tărnăuceanu and L. Tóth, *On the number of subgroups of a given exponent in a finite abelian group*, accepted for publication in Publ. Inst. Math. Beograd, arXiv:1507.00532.

Marius Tărnăuceanu
Faculty of Mathematics
"Al.I. Cuza" University
Iaşi, Romania
e-mail: tarnauc@uaic.ro